

# Anterior Esthetic Implants: Microsurgical Placement in Extraction Sockets With Immediate Provisionals

Dennis A. Shanelec, DDS

## ABSTRACT

Dentists worldwide have hoped to realize the potential for dental implants for immediate replacement of failing teeth in the maxillary esthetic zone. This article is an analysis from private practice of a case series of 100 dental implants in the anterior maxilla placed under the microscope in extraction sockets with immediate implant-supported provisionals.

The most frequent traumatic tooth injury is a fractured crown of the maxillary central incisor in the permanent dentition.<sup>1</sup> Fractured teeth frequently follow a downhill course through bonding, endodontics, posts and crowns.<sup>2</sup> Many are lost to root fracture or root resorption.<sup>3</sup> For more than 100 years, dentistry's best answer to a missing tooth in the maxillary esthetic zone has been fixed bridgework.<sup>4</sup> This was usually preceded by extraction and a removable provisional, whose other designation "the flipper" is self-descriptive of its inherent limitations. Esthetic collapse of adjacent gingival tissue and loss of buccal alveolar bone are sequela familiar to dentists restoring patients following maxillary anterior tooth loss<sup>5,6</sup> (Figure 1).

### Dental Implants and Tooth Loss

Since their introduction, dental implants have proven exceptionally predictable and successful in the edentulous and partially edentulous patient.<sup>7-10</sup> The biological and restorative improvements implants have undergone have increased their applications for single tooth replacement.<sup>11,12</sup> Dentists worldwide have hoped to realize the potential for dental implants for immediate replacement of failing teeth in the maxillary esthetic zone. This article is an analysis from private practice of a case



**Figure 1.** Ridge resorption following loss of central incisor.

series of 100 dental implants in the anterior maxilla placed under the microscope in extraction sockets with immediate implant-supported provisionals.

### Surgical Trauma and Tooth Loss

Microsurgery is a movement in medicine and dentistry toward minimally invasive alternatives to replace

**Author** / Dennis A. Shanelec, DDS, practices microsurgical periodontics and dental implants in Santa Barbara, Calif., where he also is director of the Microsurgery Training Institute.

**Disclaimer** / The author is not employed as a consultant for, nor does he receive financial support from any commercial entity.

The results described are achieved using a surgical microscope and may be statistically misleading when extrapolated to protocol not utilizing a surgical microscope.

Images conform to digital photographic ethics established by Brooks Institute of Photography: Images have not been altered other than adjustment in size, alignment, color, brightness or contrast which might have been accomplished through camera settings at the time the images were captured.

procedures that previously required extensive surgical incisions.<sup>13</sup> Using a microscope for surgery greatly enhances visual acuity and improves surgical dexterity.<sup>14</sup> Exodontia has been a traumatic procedure for centuries. Under a microscope, minimal invasive principles can be applied to tooth extraction. The aim of extraction microsurgery is to reduce trauma. Using a periosteal luxator, a tooth root can be lifted vertically from

its alveolar socket by carefully separating it from the surrounding ligament. This limits injury to the papilla and preserves natural gingival anatomy<sup>15,16</sup> (Figures 2 and 3).

The increased visibility provided under the microscope allows a surgeon to detect subtle nuances in the direction of luxation which are not apparent through normal vision, thus avoiding damaging the bone and gingival tissue.

## Implant Microsurgery

All phases of implant treatment may be performed using a microscope. Studies show that motor coordination and accuracy is generally increased when surgeons use a microscope.<sup>17</sup> Increased visual acuity, improved ergonomics, and body posture are closely related to those improvements.<sup>18</sup> In medicine, microsurgery has significantly reduced postsurgical pain in a variety of surgical disciplines.<sup>19</sup> Likewise, in endodontics, microsurgery has demonstrated measured reduction in postoperative pain.<sup>20</sup> Although no studies establish that microsurgery reduces postoperative pain following extraction or implant placement, there is strong theoretical rationale to suggest that less surgical trauma results in less pain and faster healing, and that microsurgery leads to those ends. The author's experience has shown that extraction under the microscope with implant placement in the socket followed by an immediate anatomical provisional reduces visible surgical trauma and prevents soft tissue collapse following tooth removal. Anecdotally, patients describe their discomfort level as inconsequential (Figures 4-7).

## Materials and Methods

### Patient Selection

This study comprises a case series of 100 consecutive patients in private practice requiring extraction of maxillary central incisors, lateral incisors, or cuspids. No exclusion criteria were employed. Teeth were extracted due to vertical root fracture, horizontal root fracture, root resorption or endodontic failure. One hundred implants were placed in extraction sockets and implant-supported screw-retained provisionals were delivered at the time of surgery. Preoperative and postoperative images were digitally recorded and cataloged in Extensis Portfolio image data-



**Figure 2.** Periosteal luxation of fractured root.



**Figure 3.** Periosteal extraction site. Note lack of bleeding and trauma.



**Figure 4.** Preoperative failing cuspid.



**Figure 5.** Provisional in place with screw access opening.



**Figure 6.** Provisional at time of microsurgery.



**Figure 7.** Provisional at eight weeks, ready for final restoration.

base for easy retrieval and analysis. Preoperative and postoperative radiographs were taken utilizing a Rinn film holder. These were taken without the benefit of a custom-acrylic bite plane to standardize angulation.

### *Surgical Technique*

Tooth extractions were performed using periosteal luxation. The sockets were debrided of granulation tissue and the sulcus de-epithelialized with a flame diamond. The sockets were irrigated, then filled for 30 seconds with 3 percent tetracycline solution. Lateral cutting burs were used to correct angulation between each incremental increase in twist drill size. Following completion of the osteotomies, implants were inserted utilizing a drilling unit set to 67 ncm torque and 25 rpm. All patients received NobelBiocare Mark IV 4.0 mm-diameter threaded implants with textured surfaces and 2-degree thread taper geometry.<sup>21</sup> Of 100 implants placed, 89 were

15 mm in length, nine were 18 mm in length, and two were 13 mm in length. In 81 implants, bone collected through aspiration filtration from the osteotomy preparation was rinsed with sterile water, dried and saturated with 3 percent tetracycline, then placed over any buccal thread dehiscence within the socket. Particulate bone xenograft was used to fill the remaining void between the implant and socket wall. A layer of collagen was placed over grafted bone before the implant provisional was placed. In 28 cases, preoperative buccal tissue height or thickness was judged deficient. In those cases, at the time of surgery, connective tissue was transferred from the palate into shallow subepithelial envelope incisions on the buccal. This was done to restore or maintain normal gingival height.

### *Drilling in the Extraction Socket*

Implant drilling under the microscope is a revealing experience. The

socket appears as large as a room with the apex and walls clearly visible. Drilling in extraction sites requires a different set of skills than drilling in edentulous sites.<sup>22</sup> The most favorable bone in the anterior maxilla lays to the palatal and apical of the socket<sup>23</sup> (Figures 8 and 9).

For this reason, drilling must be done at an oblique angle to the socket wall. Twist drills are not designed for this purpose. They track in the direction of less dense bone and into the open socket. Unless the osteotomy site is redirected with lateral cutting burs before each incremental increase in twist drill size, the implant angulation and position will invariable move buccally. With the magnification and lighting a microscope provides, drilling in the lateral socket wall can be accomplished for stable and accurate esthetic positioning of the implant in the socket.

### *Immediate Implant Provisional*

To preserve natural esthetics, an implant provisional must emerge from the surrounding gingiva exactly like the extracted tooth.<sup>24,25</sup> There is no margin for error. Creating a provisional crown begins before the tooth is removed.<sup>26</sup> A clear silicone impression is made, capturing the dento-gingival junction. A light-cured composite resin duplicate of the tooth is fabricated from the impression (Figures 10-12).

The duplicate tooth is trimmed to the exact location of the dento-gingival



**Figure 8.** Implant properly placed in lingual side of socket. Note immediate tissue collapse on the buccal without provisional support.



**Figure 9.** Lateral cutting burs for osteotomy preparation in the lateral wall of the socket.



**Figure 10.** Blank replica of failing tooth.



**Figure 11.** Replica tooth hollowed and shaped. The rough surface facilitates adding flowable composite to create ideal tissue support.



**Figure 12.** Polished and glazed provisional with proper subgingival profile.



**Figure 13.** Lateral incisor removed micro Surgically with implant in place.



**Figure 14.** Opaqued titanium screw-retained abutment on implant.



**Figure 15.** Replica tooth fitted to abutment.



**Figure 16.** Replica tooth bonded to the temporary abutment with light-cured composite.



**Figure 17.** Final finished and polished provisional crown.



**Figure 18.** Provisional crown at time of implant microsurgery.



**Figure 19.** Provisional crown at eight weeks, ready for final restoration.

junction. The replica tooth is hollowed for luting to a titanium screw-retained temporary abutment. The outer surface of the titanium abutment is opaqued to provide accurate color match. The temporary titanium abutment is placed on the implant and the duplicate crown filled with light-cured composite then joined to the abutment. (Figures 13 and 14).

This provisional crown is removed

utilizing a screw access opening in the incisal third. The subgingival profile is individually shaped for each patient at the time of surgery.<sup>27</sup> Voids and rough edges are eliminated, and the provisional carefully contoured to support the gingival tissue. As a final step, it is polished and glazed (Figures 15-18).

Using light-cured composite assures that no free monomer is present to irritate tissue or bone. The machined titanium provisional abutment reduces the possibility of the provisional loosening. In this study, of the 100 provisionals placed, screw loosening occurred in one patient (Figure 19).

### *Provisional Occlusion*

Early loading bone forces are controlled in multiple implant immediate loading cases through splinting.<sup>28</sup> Early loading bone forces in single implant cases are controlled by eliminating centric and lateral occlusal contact. Symmetrical and light mesial and distal contacts are established and the provisional is taken completely out of centric and lateral occlusal contact using green occlusal indicator wax. This protocol allows patients to leave the dental office with a non-loaded esthetic provisional tooth securely anchored to the dental implant.

### *Final Esthetic Restoration*

An immediate provisional crown assures that patients are never without a natural-looking tooth. Because the gingiva is never unsupported, natural tissue height and contour can be preserved. The subgingival emergence profile of the provisional must be accurately registered and transferred to the ceramist.<sup>29-31</sup> This transfer of subgingival anatomy is accomplished through an impression made of the gingival third of the provisional crown with an implant analog attached. The impression is used to create a custom impression transfer coping which duplicates

the subgingival emergence profile of the provisional<sup>32-34</sup> (Figures 20-23).

After final impressions with the custom impression transfer coping, computer-assisted scanning and machining then creates a final Procera Zirconia ceramic abutment and ceramic crown<sup>35,36</sup> (Figures 24 and 25).

This process assures that the final abutment exactly matches both the original tooth shape and the provisional emergence profile. Working together as a team, surgeons, restorative dentists, and ceramists can combine their skill and knowledge to create a tooth in harmony with adjacent gingival anatomy and the overall appearance of face and smile (Figures 26-29).

For many patients, traumatic anterior tooth loss is their first exposure to implant dentistry. In the author's experience, patients are pleased with the simplicity of a technique which does not involve major incisions, suturing, raising of mucogingival flaps or the need for multiple surgical procedures. They frequently comment on the "sturdiness" of their provisional and lack of pain associated with the procedure (Figure 30).


### Clinical Results

Using the described protocol, 100 implants were placed over a 36-month period. The average follow-up time was 18 months. Of 100 implants placed, 98 implants were approved for restoration at eight weeks and successfully restored. Two implants failed to integrate before restoration. These were removed and replaced, then provisionalized with bonded pontic provisionals prior to restoration. The clinical success rate for implants placed using this protocol was 98 percent based on the following parameters:

- Absence of clinical inflammation, infection or bleeding on probing
- Absence of mobility, pain or sensitivity of the implant or surrounding tissues



**Figure 20.** Impression of provisional crown is taken before it is placed on the implant.



**Figure 21.** Registration of the provisional crown emergence profile.



**Figure 22.** Clear acrylic reproduces provisional emergence profile.



**Figure 23.** Custom impression transfer coping polished and glazed.



**Figure 24.** Zirconia all-ceramic implant abutment.



**Figure 25.** Procera all-ceramic crown

- Soft tissue sulcus less than 1 mm beyond the implant platform
- Ability to withstand rotational torque of 45 ncm at abutment insertion
- Radiographic evidence of bone to within the topmost thread of the implant
- The implant is restored and remains in clinical function

### Discussion

Dental implants originally followed a protocol which required lengthy periods

of undisturbed healing before loading, typically three and six months.<sup>37-39</sup> Immediately loading of full arch splinted implants is now accepted therapy.<sup>40-43</sup> Immediate loading of single tooth implants has also proven successful.<sup>44-49</sup> Combined data from six studies, comprising 287 single tooth implants immediately loaded showed a 96.7 percent survival rate.<sup>50</sup> Seven studies have examined the outcome of implants placed in extraction sockets of the anterior maxilla

with immediate provisionals.<sup>51-57</sup> Combined data from these studies comprised 190 implants and showed a 95.9 percent survival rate.<sup>58</sup> The successful outcomes described by these investigators have proven controversial. Reviewing these seven studies, Ganeles and Wismeijer stated, "It should be recognized that, with few exceptions, these conclusions may be misleading statistical phenomena of the authors, as most publications were written by exceptionally experienced, highly skilled practitioners working under tightly controlled clinical conditions on a relatively small, statistically inconclusive number of implants and patients."

Such ad hominem arguments marginalizing obviously successful results reflect the frustration experienced by many clinicians who attempt immediate provisionals on implants placed in extraction sockets. However, our current study confirms a high clinical success rate of implants placed microsurgically in extraction sockets of the anterior maxilla with immediate provisionals. Comprising 100 consecutive implants, it is the largest study to date examining this protocol. The operator skill requirements and technical demands of the protocol are admittedly high. It is the author's opinion that optical magnification afforded by the microscope increases the precision of placement and initial stability during implant microsurgery. In addition, the minimal invasiveness and reduced surgical trauma of microsurgically placed dental implants may contribute to rapid healing, lessened morbidity and successful osseointegration.

Clinical success describes the basic survival, health, comfort, and function of dental implants but does not define esthetic success. In implant dentistry, evidence-based statements regarding esthetic procedures are difficult to generate. Most studies focus on implant survival. Scientific evidence of estheti-



**Figure 26.** Preoperative X-ray of failing central incisor.



**Figure 27.** Preoperative central incisor.



**Figure 28.** Postoperative implant at 44 months.



**Figure 29.** Postoperative X-ray at 44 months.

cally reproducible parameters is rare. The stability of soft tissue esthetics around single tooth implants has been studied. Significant regeneration of mesial and distal papilla was shown after a follow-up period of 1.5 years.<sup>59</sup> On the other hand, soft tissue buccal recession of 0.6 mm was also shown after one year.<sup>60</sup> This has led to a consensus that a provisional restoration with adequate emergence profile should be used to guide and shape the peri-implant tissue prior to final restoration.<sup>61</sup> Priest proposed an esthetic analysis based on soft tissue gingival height around dental implants. In his analysis, the positions of midfacial gingival tissue margin and the heights of mesial and distal papillae were numerically compared using fixed reference points on preoperative and postoperative photographs. Our current study utilized a variation of Priest's analysis as a

subjective measure of esthetic outcome on a visual analog scale. Based on before and after digital photographs, the gingival anatomy remained unchanged or improved in 95 patients. Five patients required postoperative connective tissue grafts to bring the gingival zenith to preoperative levels. A more comprehensive numerical digital analysis of this case series will be forthcoming in future publications.

Belser, Buser and Higginbottom defined an esthetic implant crown as one in harmony with the perioral facial structures. In addition, the soft tissues, including, height, volume, color, texture and contours, should be in harmony with the surrounding teeth. Finally, the restoration should imitate the natural appearance of the missing tooth.<sup>62</sup> This study shows that microsurgery can be utilized for implant placement in extraction sockets with a high degree of



**Figure 30.** Smile line esthetics with implant and crown (Restoration courtesy Kathleen McClintock, DDS).

clinical success. When provisionals are placed at the time of surgery, the gingival tissues are supported and undergo little change in anatomy. This protocol offers an opportunity for implant therapy with less morbidity and highly esthetic outcomes. This translates into increased patient acceptance and satisfaction.<sup>63</sup> Dentistry will see increasing use of the microscope in clinical practice, including applications for extraction, implant placement and restoration.<sup>64</sup> Microscopy has the potential to advance dentistry from an era of traumatic tooth loss to one of exact and seamless replacement of a failing anterior tooth with an esthetic implant-supported crown. CDA

**References** / 1. Bastone EB, Freer TJ, McNamara JR, Epidemiology of dental trauma: a review of the literature. *Aust Dent J* 45(1):2-9, March 2000.

2. Al-Jundi SH, Dental emergencies presenting to a dental teaching hospital due to complications from traumatic dental injuries. *Dent Traumatol* 18(4):181-5, August 2002.

3. Andreasen FM, Transient root resorption after dental trauma: the clinician's dilemma *J Esthet Restor Dent* 15(2):80-92, 2003.

4. Studer S, Pietrobon N, Wohlwend A, Maxillary anterior single-tooth replacement: a comparison of three treatment modalities. *Pract Periodontics Aesthet Dent* 6(1):51-60, January-February 1994.

5. Small BW, Achieving and maintaining periodontal health and esthetics following the extraction of a central incisor. *Gen Dent* 51(5):396-8, September-October 2003.

6. Nicopoulou-Karayianni K, Bragger U, Lang NP, Patterns of periodontal destruction associated with incomplete root fractures. *Dentomaxillofac Radiol* 26(6):321-6, November 1997.

7. Schroeder A, Pohler O, Sutter E,

Gewebsreaktion auf ein Titan-hohlzylinder-implantat mit Titan-Spritzschichtober-fläche. *Monatsschr Zahnheilkd* 86:713-27, 1976.

8. Adell R, Lekholm, U, Rockler B, Brånemark PI, A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 10:387-416, 1981.

9. Adell R, Erikson B, Lekholm U, Brånemark PI, A long-term follow-up study of osseointegrated implants in treatment of the totally edentulous jaws. *Int J Oral Maxillofac Implants* 5:347-58, 1990.

10. Jemt T, Petterson P, A three-year follow-up study on single implant treatment. *J Dent* 21:203-8, 1993.

11. Linquist LW, Carlsson GE, Jemt T, A prospective 15-year follow-up study of mandibular fixed prosthesis supported by osseointegrated implants. *Clin Oral Implants Res* 7:329-36, 1996.

12. Widmark G, Friberg B, Johansson B, Sindet-Pederson S, Taylor A, Mk III: a third-generation of the self-tapping Brånemark System implant, including the new Stargrip internal grip design. A one-year prospective four-center study. *Clin Implant Dent Relat Res* 3:5(4):273-9, 2003.

13. Serafin D, Microsurgery: Past Present and Future. *Plastic and Reconstr Surg* 66:781-5, 1980.

14. Stephans JA, Taylor A, The effect of visual feedback on physiological tremor, *EEG Clin Neurophysiol* 34:457-64, 1974.

15. Handtmann S, Lindemann W, Application possibilities of the periosteome for various technics in periodontal surgery. *Dtsch Zahnärztl Z* 40(7):745-8, July 1985.

16. Thomson PJ, Minimising trauma in dental extractions: the use of the periosteome. *Br Dent J* 7:172(5):179, March 1992.

17. Harris H, Mackensen, G, Ocular surgery under the microscope, Chicago, Yearbook Medical Publishers, Inc., 1987

18. Leknius C, Geissberger, M, The effect of magnification on the performance of fixed prosthodontic procedures. *J Calif Dent Assoc* 23(12):66-70, 1995.

19. Pecora G, Andreana S, Operating microscope in endodontic surgery. *Oral Surg Med Pathol* 75:751, 1993.

20. Daniel RK, Microsurgery: Through the Looking Glass, *N Engl J Med* 300: 1251-8, 1979.

21. O'Sullivan D, Sennerby L, Meredith N, Influence of implant taper on primary and secondary stability of osseointegrated titanium implants *Clin Oral Implants Res* 15:474-80, 2004.

22. Campelo LD, Dominguez Camara J, Flapless implant surgery: a 10-year clinical retrospective analysis. *Int J Oral Maxillofac Implants* 17(2):271-6, 2002.

23. Hartman G, Cochran D, Initial implant position determines the magnitude of crestal bone remodeling. *J Periodontol* 75(40):572-7, 2004.

24. Saadoun AP, Le Gall M, Implant positioning for periodontal, functional and aesthetic result. *Pract Periodontics Aesthet Dent* 4:43-54, 1992.

25. Saadoun AP, Immediate implant placement and temporization in extraction and healing sites. *Compend Contin Educ Dent* 23(4):309-324, April 2002.

26. Macintosh DC, Sutherland M, Method for developing an optimal profile using heat-polymerized provisional restorations for single-tooth implant-supported restorations. *J Prosthet Dent* 91(3):289-92, March 2004.

27. King KO, Implant abutment emergence profile: key to esthetics. *J Oral Implantol* 22(1):27-30, 1996.

28. Bain WA, Weisgold AS, Customized emer-

gence profile in the implant crown – a new technique. *Compend Contin Educ Dent* 18(1):41-5; quiz 46, January 1997.

29. Misch CE, Wang HL, Misch CM, Sharawy M, Lemons J, Judy KW, Rationale for the application of immediate load in implant dentistry: Part I. *Implant Dent* 13(3):207-17, 2004.

30. Breeding LC, Dixon DL, Transfer of gingival contours to the master cast. *J Prosthet Dent* 70:341-3, 1993.

31. Hinds KF, Custom impression coping for an exact registration of the healed tissue in the esthetic implant restoration. *Int J Periodontics Restorative Dent* 17(6):584-91, December 1997.

32. Buskin R, Salinas TJ, Transferring emergence profile created from the provisional to the definitive restoration. *Pract Periodontics Aesthet Dent* 10(9):1171-9, November-December 1998.

33. Attard N, Barzilay I, A modified impression technique for accurate registration of peri-implant soft tissues. *J Can Dent Assoc* 69(2):80-3, February 2003.

34. Touati B, Guez G, Saadoun A, Aesthetic soft tissue integration and optimized emergence profile: provisionalization and customized impression coping. *Pract Periodontics Aesthet Dent* 11(3):305-14, April 1999.

35. Davarpanah M, Martinez H, Celletti R, Tecucianu JF, Three-stage approach to aesthetic implant restoration: emergence profile concept. *Pract Proced Aesthet Dent* 13(9):761-7, November-December 2001.

36. Yildirim M, Edelhoff, D, Hanisch O, Spiekermann H, Ceramic abutments – a new era in achieving optimal esthetics in implant dentistry. *Int J Periodontics Restorative Dent* 20(1):81-91, February 2000.

37. Brånemark PI, Hansson BO, Adell R, et al, Osseointegrated implants in the treatment of the edentulous jaw. *Scand J Plast Reconstr Surg* 16:1-132, 1977.

38. Adell R, Lekholm U, Rockler B, Brånemark PI, A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 10:387-416, 1981.

39. Buser D, Merieske-Stern R, Bernard JP, et al, Long-term evaluation of non-submerged ITI implants. Part I: Eight-year life table analysis of a prospective multicenter study with 2,359 implants. *Clin Oral Implants Res* 8:161-72, 1997.

40. Rocuzzo M, Bunino M, Prioglio F, Bianchi SD, Early loading of sand blasted and acid-etched (SLA) implants: a prospective split-mouth comparative study. *Clin Oral Implants Res* 12:572-8, 2001.

41. Rocuzzo M, Wilson T, A prospective study evaluating a protocol for six weeks' loading of SLA implants in the posterior maxilla: one-year results. *Clin Oral Implants Res* 13:502-7, 2002.

42. Testori T, Del Fabrio M, Feldman S, et al, A multicenter prospective evaluation of two-month loaded Osseotite implants placed in the posterior jaws: Three-year follow-up results. *Clin Oral Implants Res* 13:154-61, 2002.

43. Bogaerde IV, Pedretti G, Dellacassa P, Mozzati M, Rangert B, Early function of splinted implants in maxillas and posterior mandibles using Brånemark system machined-surface implants: an 18-month prospective clinical multicenter study. *Clin Implant Dent Relat Res* 5(suppl 1):21-8, 2003.

44. Malo P, Rangert B, Dvarsater L, Immediate function of Brånemark implants in the esthetic zone: a retrospective clinical study with six months to four years of follow up. *Clin Implant Dent Relat Res* 2:138-46, 2000.

45. Hui E, Chow J, Li D, Liu J, Wat P, Law H,

The Dental Record new

Immediate provisionalization for single tooth implant replacement with Brånemark system: preliminary report. *Clin Implant Dent Relat Res* 3:79-86, 2001.

46. Chaushu G, Chaushu S, Tzohar A, Dayan D, Immediate loading of single tooth implants: immediate versus nonimmediate implantation. *Int J Oral Maxillofac Implants* 16:267-72, 2001.

47. Malo P, Friberg B, Polizzi G, Gualini F, Vighagen T, Rangert B, Immediate and Early Function of Brånemark System implants placed in the esthetic zone: a 1-year prospective clinical multicenter study. *Clin Implant Dent Relat Res* 5(suppl 1):37-46, 2003.

48. Glauser R, Lundgren AK, Gottlw, J, et al, Immediate occlusal loading of Brånemark TiUnite implants placed predominantly in soft bone: 1-year results of a prospective clinical study. *Clin Implant Dent Relat Res* 5(suppl 1):47-55, 2003.

49. Cannizzaro G, Leone M, Restoration of partially edentulous patients using dental implants with a microtextured surface: a prospective comparison of delayed and immediate full occlusal loading. *Int J Oral Maxillofac Implants* 18:512-22, 2003.

50. Ganeles J, Wismeijer D, Early and immediately restored and loaded dental implants for single-tooth and partial-arch applications. *Int J Oral Maxillofac Implants* 19(supplement):92-102, 2004.

51. Kan YK, Rungcharassaeng K, Lozada J, Immediate placement and provisionalization of maxillary anterior single implants: One-year prospective study. *Int J Oral Maxillofac Implants* 30:18:31-9, 2000.

52. Wohrle PS, Single-tooth replacement in the esthetic zone with immediate provisionalization: fourteen consecutive case reports. *Pract Periodontics Aesthet Dent* 10:1107-14, 1998.

53. Malo P, Rangert B, Dvarsater L, Immediate function of Brånemark implants in the esthetic zone: a retrospective clinical study with 6 months to 4 years of follow-up. *Clin Implant Dent Relat Res* 2:138-46, 2000.

54. Hui E, Chow J, Li D, Liu J, Wat P, Law H, Immediate provisionalization for single tooth implant replacement with Brånemark system: preliminary report. *Clin Implant Dent Relat Res* 3:79-86, 2001.

55. Chaushu G, Chaushu S, Tzohar A, Dayan D, Immediate loading of single tooth implants: immediate versus non-immediate implantation. *Int J Oral Maxillofac Implants* 16:267-72, 2001.

56. Malo P, Friberg B, Polizzi G, Gualini F, Vighagen T, Rangert B, Immediate and Early Function of Brånemark System implants placed in the esthetic zone: a One-year prospective clinical multicenter study. *Clin Implant Dent Relat Res* 5(suppl 1):37-46, 2003.

57. Kan YK, Rungcharassaeng K, Lozada J, Immediate placement and provisionalization of maxillary anterior single implants: One-year prospective study. *Int J Oral Maxillofac Implants* 30:18:31-9, 2000.

58. Grunder U, Stability of the mucosal topography around single tooth implants and adjacent teeth. *Int J Periodontics Restorative Dent* 207:11-7, 2000.

59. Jemt T, Regeneration of gingival papilla after single implant treatment. *Int J Periodontics Restorative Dent* 17:326-33, 1997.

60. Belser U, Buser D, Higginbottom F, Consensus statements and recommended clinical procedures regarding esthetics in implant dentistry. *Int J Oral Maxillofac Implants* 19(supplement):73-4, 2004.

61. Belser U, Schmid B, Higginbottom F, Buser

D, Outcome analysis of implant restoration located in the anterior maxilla: a review of the recent literature. *Int J Oral Maxillofac Implants* 19(supplement):30-42, 2004.

62. Priest G, Predictability of soft tissue form around single-tooth implant restorations. *Int J Periodontics Restorative Dent* 23:19-27, 2003.

63. Schropp L, Flemming I, Kostopoulos L, Wenzel A, Patient experience of, and satisfaction with delayed-immediate vs. delayed single-tooth implant placement. *Clin Oral Implants Res* 15:490-7, 2004.

64. Shanelec, D, Tibbetts, L, Recent Advances in Surgical Technology, Clinical Periodontology 8th edition, W. B. Saunders 2004.

**To request a printed copy of this article, please contact / Dennis A Shanelec, DDS, director, Microsurgery Training Institute, 1515 State St., Suite 1, Santa Barbara, CA 93101.**